

**MAPPING PM_{2.5} POLLUTION ZONES IN NIGERIA USING HIERARCHICAL CLUSTERING*****¹Emmanuel Yohanna, ¹Jasini Waida, ²Yusuf Emmanuel Yila**¹Department of Physics, Borno State University, Maiduguri, Borno State.²Chemistry Department, Federal University of Agriculture Markurdi, Benue State.*Corresponding authors' email: wabuemma@gmail.com**ABSTRACT**

The persistent challenges in Nigeria's air-quality monitoring network have limited large-scale spatial assessments of particulate matter. This study conducts a national-scale clustering analysis of PM_{2.5} using agglomerative hierarchical clustering which does not require a pre-specified number of clusters and shows hierarchical linkages across monitoring sites. Ward's method was applied to daily PM_{2.5} data from 15 monitoring stations over the period from 2019 to 2023, using Euclidean distance to assess the similarity measure. Three distinct clusters were identified: High Pollution Regions (HPR) with an average PM_{2.5} concentration of 61.7 $\mu\text{g m}^{-3}$, Medium Pollution Regions (MPR) at 46.7 $\mu\text{g m}^{-3}$, and Low Pollution Regions (LPR) at 36.0 $\mu\text{g m}^{-3}$. The 2019 haze episode produced peak concentrations of 197.7 $\mu\text{g m}^{-3}$ in HPR, leading to a hazardous Air Quality Index (AQI) of 273. Cluster distributions demonstrated a significant correlation with land-use characteristics. HPR stations were primarily situated in industrial zones, MPR in densely built urban corridors, and LPR in coastal peri-urban regions. The resulting novel spatial stratification offers substantial evidence for developing targeted and regionally adaptive air quality management strategies across Nigeria.

Keywords: Agglomerative Hierarchical Cluster, Air Pollution, Air Quality Index, Nigeria, PM_{2.5}**INTRODUCTION**

Air pollution is a significant environmental risk to global health, associated with approximately seven million premature deaths annually (Manisalidis et al., 2020; World Health Organization, 2021; Khomenko et al., 2021). Though, fine particulate matter (PM_{2.5}) can penetrate deep into the lungs and enter the bloodstream, significantly contributing to the burden of cardiopulmonary morbidity and mortality (Benabed & Boulbair, 2022; Garcia et al., 2023). But, PM_{2.5} pollution is prevalent in sub-Saharan Africa; however, significant areas of the region are inadequately monitored, leading to considerable deficiencies in exposure assessment and risk estimation (Glenn et al., 2022; World Health Organization, 2022; Lawal & Muhammed, 2022).

Nigeria exemplifies this challenge distinctly. The country, with a population surpassing 200 million, confronts increasing pollution challenges due to rapid urbanization, growing industrial activities, congested transportation systems, and frequent biomass burning (Busolo & Njabira, 2022; Almsatar, 2020). National PM_{2.5} levels are approximately five times higher than the WHO guideline (Ezeh et al., 2019; Wambebe & Duan, 2020). However, major cities, including Lagos, Kano, Abuja, and Port Harcourt, frequently encounter unhealthy to hazardous concentrations, especially during the dry season (Sunday, 2022). Spatial heterogeneity is evident across the country's ecological zones, influenced by industrial distribution, vehicle density, meteorological variability, and transboundary haze transport (Ogwu et al., 2024; Abulude et al., 2024). Despite the magnitude of these challenges, Nigeria's monitoring capacity is insufficient; a 2024 audit reveals that the majority of stations are either non-functional or exhibit limited reliability (Omokaro et al., 2025).

The emission landscape highlights the intricacies of exposure. Vehicular emissions are prevalent in urban environments, exacerbated by the importation of outdated and inadequately regulated vehicles (Singh et al., 2023; Oluwakoya, 2024), while, industrial sources, especially within the oil-producing corridor of the Niger Delta, significantly contribute to elevated levels of PM_{2.5} and gas-phase pollutants (Meo et al.,

2024; World Health Organization, 2021). But, biomass combustion and the widespread utilization of diesel and petrol generators during power outages contribute to persistent emissions in rural and peri-urban communities. Seasonal dynamics, particularly during the Harmattan period, increase PM_{2.5} levels by 20–60% due to the advection of Saharan dust across the region (Léon et al., 2021).

Also, under conditions characterized by spatial diversity and limited data, it is crucial to employ analytical methods that can effectively extract robust spatial structures from sparse monitoring networks. Agglomerative Hierarchical Clustering (AHC) presents several advantages in this context. It eliminates the need for pre-defining the number of clusters, elucidates hierarchical relationships, and demonstrates effectiveness in heterogeneous pollution environments (Paparrizos et al., 2024; Mitchell & Bala, 2024). Meanwhile, research conducted in Malaysia, India, and Europe indicates that Agglomerative Hierarchical Clustering (AHC) employing Ward's method and Euclidean distance successfully categorizes regions into high, medium, and low pollution classifications, thereby enhancing the precision of regulatory and public health interventions (Rahman et al., 2022; Azizan et al., 2023). The classification has been validated with high accuracy through discriminant analysis (Azizan et al., 2023).

Despite the relevance of this approach, no nationwide AHC-based assessment of PM_{2.5} spatial patterns has been conducted in Nigeria, leaving a critical evidence gap in the country's air-quality management framework. This study addresses that gap by applying hierarchical clustering to 5 years of PM_{2.5} records from Nigeria's available monitoring stations. This work is guided by three research questions:

- Can Ward's agglomerative hierarchical clustering delineate coherent PM_{2.5} pollution regions across Nigeria?
- How do PM_{2.5} levels vary temporally across these regions between 2019 and 2023?
- What do associated AQI values reveal about potential health risks across the identified pollution zones?

By answering these questions, the study provides the first national-scale, cluster-based characterization of PM_{2.5} in Nigeria and delivers policy-relevant insights for strengthening air-quality governance in one of Africa’s most polluted and data-constrained environments.

MATERIALS AND METHODS

Study Area

Figure 1 shows Nigeria map which spans 923,768 km² in West Africa, between latitudes 4°N and 14°N and longitudes

3°E and 14°E (Adeleye et al., 2021), with a tropical climate with wet and dry seasons. The rainy season is April to October, while the dry season is November to March. Sufiyan et al. (2020) reported that the Harmattan wind blows dusty Sahara Desert air during the dry season, affecting air quality. Nigeria is divided into six geopolitical regions, South South, South West, South East, North Central, North West, and North East, each characterized by unique economic activities, varying population densities, and specific sources of pollution (Elechi et al., 2023).



Figure 1: Map of Nigeria Showing the Locations. Britannica, Inc. (2025)

Air Quality Monitoring Stations

This study utilized data gathered from 15 air quality monitoring stations that are strategically located throughout Nigeria’s six geopolitical zones, as shown in Table 1. The selection of stations was determined by factors such as geographical representation and classification into urban, industrial, and suburban categories. The stations include seven (7) urban stations positioned in central urban areas

characterized by significant traffic volume, five (5) suburban stations located in residential neighborhoods characterized by moderate development, and three (3) industrial stations located in proximity to oil refineries, manufacturing plants, and industrial zones. The coordinates of the stations varied between 4.82°N and 12.00°N latitude, as well as 3.35°E and 13.08°E longitude, providing extensive spatial coverage.

Table 1: Air Quality Monitoring Stations in Nigeria

Station Code	Station Name	State	Zone	Location Type	Coordinates
NIG01	Lagos-Ikeja	Lagos	Southwest	Urban	6.5977°N, 3.3464°E
NIG02	Lagos-Lekki	Lagos	Southwest	Suburban	6.4437°N, 3.4700°E
NIG03	Abuja-Municipal	FCT Abuja	North Central	Urban	9.0574°N, 7.4898°E
NIG04	Abuja-Asokoro	FCT Abuja	North Central	Suburban	9.0365°N, 7.5339°E
NIG05	Kano-City	Kano	Northwest	Urban	12.0022°N, 8.5920°E
NIG06	Kano-Industrial	Kano	Northwest	Industrial	11.9716°N, 8.5378°E
NIG07	Port Harcourt-Trans Amadi	Rivers	South South	Industrial	4.8156°N, 7.0498°E
NIG08	Port Harcourt-Rumuola	Rivers	South South	Urban	4.8891°N, 7.0229°E
NIG09	Benin City-Central	Edo	South South	Urban	6.3381°N, 5.6157°E
NIG10	Benin City-Airport Road	Edo	South South	Suburban	6.3176°N, 5.6347°E

Station Code	Station Name	State	Zone	Location Type	Coordinates
NIG11	Enugu-Coal Camp	Enugu	Southeast	Suburban	6.4398°N, 7.4951°E
NIG12	Ibadan-Bodija	Oyo	Southwest	Urban	7.3876°N, 3.9093°E
NIG13	Warri-Effurun	Delta	South South	Urban	5.5378°N, 5.7821°E
NIG14	Maiduguri-GRA	Borno	Northeast	Urban	11.8469°N, 13.0820°E
NIG15	Calabar-Marian	Cross River	South South	Suburban	4.9517°N, 8.3417°E

PM_{2.5} Measurement and Data Collection

PM_{2.5} concentrations were measured through a network of stations equipped with Tapered Element Oscillating Microbalance (TEOM 1405/1400a) and Beta Attenuation Monitors (BAM-1020), both recognized as Federal Equivalent Methods by the U.S. EPA (Magi et al., 2020; Jin et al., 2019). TEOM units were primarily utilized at southern and coastal locations, equipped with humidity control modules to reduce high-moisture interference. In contrast, BAM monitors were positioned at northern and inland stations to ensure stable performance in drier environments (Jin et al., 2019).

Hourly measurements were compiled into 24-hour averages from 1 January 2019 to 31 December 2023, yielding 1,826 daily observations per station (total n = 27,390). Outlier detection utilized the interquartile range (IQR) method for the daily series of each station. Among the flagged values, 147 observations (0.54% of the dataset) were initially identified as potential outliers; 112 were confirmed as instrument or power-related anomalies and subsequently removed, while 35 were retained following cross-validation with calibration and meteorological logs. Data gaps, representing less than 5% of total observations, were addressed through linear interpolation. However, longer periods of missing data were omitted from further analyses.

Agglomerative Hierarchical Cluster Analysis

Agglomerative Hierarchical Clustering (AHC) was employed to identify spatial groupings of stations based on similarities in PM_{2.5} levels (Azizan et al., 2023; Rahman et al., 2022). This was conducted to ascertain the spatial distribution of the stations. Maldonado-Salguero et al. (2022) indicate that the method commenced with each station operating as an individual cluster. Subsequently, an iterative process was employed to merge the clusters. Ward's linkage was selected due to its ability to minimize variation within clusters, resulting in compact groups that are distinctly separated from each other (El-Araby et al., 2023). Metrics, including median, percentiles, and variability measures, provide various insights into air quality. However, long-term mean PM_{2.5} is preferred for clustering due to its representation of chronic exposure, which is the main contributor to cardiovascular and respiratory risks, and its compliance with WHO and U.S. EPA standards (Zhou et al., 2025; Roy et al., 2025; Jalali et al., 2021). The mean, unlike medians or short-term variability, mitigates the impact of episodic extremes and offers a stable,

comparable metric across stations, thereby serving as an optimal tool for detecting enduring spatial patterns within Nigeria's PM_{2.5} network as illustrated in Eq. 1.

$$\Delta S = \frac{p_i q_j}{p_i + q_j} \|q_i - q_j\|^2 \tag{1}$$

where p_i and p_j are the number of observations in clusters i and j, and q_i and q_j are the cluster centroids.

Distance Metric

Euclidean distance was employed as illustrated in Eq. 2 (Balaji et al., 2022) as the dissimilarity measure, calculated as:

$$E_{ij}^2 = \sum_{k=1}^n (d_{ik} - d_{jk})^2 \tag{2}$$

In this context, E²_{ij} refers to the squared Euclidean distance between stations i and j, while d_{ik} denotes the mean PM_{2.5} concentrations at station i. Additionally, n represents the number of variables, which in this instance is n = 1, indicating the mean PM_{2.5}. (Fu et al., 2020). The linkage distance was standardized using the formula (E_{link}/E_{max}) × 100, with E_{link} representing the distance at which clusters converge and E_{max} indicating the maximum distance within the dataset. This normalization enhances the understanding of the y-axis in the dendrogram used in the results analysis. The AHC analysis utilized the SciPy library in Python. The process included determining the average PM_{2.5} concentrations for each station from 2019 to 2023. Calculating the Euclidean distances among all stations and utilizing Ward's linkage algorithm to create a dendrogram visualization and determine the optimal height for cutting the dendrogram to yield three distinct clusters. Assigning cluster labels of HPR, MPR, and LPR according to the average PM_{2.5} values was also implemented.

Air Quality Index

The Air Quality Index (AQI) was determined utilizing the updated breakpoints for PM_{2.5} as provided by US EPA PM_{2.5} AQI (2024) and Horn & Dasgupta (2024), as seen in Table 2. AQI value formula is stated as Eq. 3 (Pyae, 2021),

$$I_p = \frac{I_{Hi} - I_{Lo}}{BP_{Hi} - BP_{Lo}} (C_p - BP_{Lo}) + I_{Lo} \tag{3}$$

where:

I_p = AQI value for pollutant p (PM_{2.5}), C_p = Truncated concentration of PM_{2.5} (to 1 decimal place), BP_{Hi} = Concentration breakpoint ≥ C_p, BP_{Lo} = Concentration breakpoint ≤ C_p, I_{Hi} = AQI value corresponding to BP_{Hi}, I_{Lo} = AQI value corresponding to BP_{Lo}

Table 2: US EPA PM_{2.5} AQI Range (2024)

AQI Category	AQI Range	PM _{2.5} Range (µg/m ³)	Health Implications
Good	0-50	0.0-9.0	Air quality is satisfactory
Moderate	51-100	9.1-35.4	Acceptable quality
Unhealthy for Sensitive Groups	101-150	35.5-55.4	Sensitive groups may experience effects

Unhealthy	151-200	55.5-125.4	Everyone may experience health effects
Very Unhealthy	201-300	125.5-225.4	Health alert conditions
Hazardous	301+	225.5+	Emergency conditions

Data visualization analyses and visualizations were conducted using Python (version 3.13.5). The workflow including data preprocessing, normalization, clustering and performance assessment and evaluation using libraries such as scikit-learn, pandas, NumPy, and matplotlib. Descriptive statistics such as mean, median, standard deviation, skewness, and kurtosis. Positive skewness values suggest right-tailed distributions typical of pollution events (Rahman et al., 2022). Temporal trends were examined through the aggregation of monthly and annual data.

RESULTS AND DISCUSSION

Spatial Distribution of PM_{2.5} Concentrations

Over a five-year period, PM_{2.5} concentrations at 15 Nigerian monitoring stations ranged from 33.7 to 67.2 µg m⁻³, exceeding the WHO annual recommendation of 5 µg m⁻³ by factors of 6.7 to 13.4, as shown in table 3. Concentrations were highest in industrialized urban centers, including Kano-Industrial (67.2 µg m⁻³), Port Harcourt-Trans Amadi (59.9 µg

m⁻³), and Kano-City (57.8 µg m⁻³), which are emission hotspots in Nigeria's manufacturing and petroleum refining areas. Lower but significantly increased values were seen at semi-rural sites such as Calabar Marian (33.7 µg m⁻³), Enugu Coal Camp (35.9 µg m⁻³), and Abuja Asokoro (38.4 µg m⁻³). Positive skewness (1.07–1.91) and leptokurtic distributions (kurtosis: 2.42–8.99) indicated rare high-pollution episodes over baseline concentrations. Warri-Effurun had the highest skewness at 1.91, reflecting Nigeria's oil and gas sector's interrupted industrial emissions. Distributional features show that long-term averaging in air quality management is insufficient and suggest that policy frameworks should account for episodic pollution spikes. Nigeria's exceedance of WHO guidance, compared to sub-Saharan African baselines where many cities report similar or elevated PM_{2.5} levels, highlights regional air quality issues caused by rapid industrialization, insufficient emissions control, and biomass burning.

Table 3: Descriptive Statistics of PM_{2.5} (µg/m³) by Station from 2019 to 2023

Station Code	Station Name	State	Region Type	Min	Q1	Median	Q3	Max	Mean	SD	Skewness	Kurtosis
NIG06	Kano-Industrial	Kano	HPR	24.08	49.74	61.78	81.76	241.32	67.20	23.84	1.39	4.12
NIG07	PortHarcourt-TransAmadi	Rivers	HPR	20.41	45.08	55.47	73.35	197.61	59.97	20.08	1.18	3.25
NIG05	Kano-City	Kano	HPR	18.64	43.02	53.70	69.90	228.97	57.81	19.79	1.42	5.57
NIG09	BeninCity-Central	Edo	MPR	19.89	39.76	48.88	64.17	180.72	52.58	16.98	1.17	3.93
NIG01	Lagos-Ikeja	Lagos	MPR	14.94	37.82	46.58	60.89	181.85	50.57	17.86	1.58	5.56
NIG13	Warri-Effurun	Delta	MPR	16.64	37.21	45.98	59.46	189.00	49.60	17.61	1.91	8.99
NIG08	PortHarcourt-Rumuola	Rivers	MPR	16.89	36.39	44.82	58.43	158.67	48.22	15.75	1.07	2.42
NIG03	Abuja-Municipal	FCT	MPR	13.67	34.01	41.94	55.66	182.41	46.01	16.75	1.76	7.03
NIG12	Ibadan-Bodija	Oyo	MPR	14.94	34.30	42.43	55.64	166.46	45.63	15.41	1.33	4.82
NIG10	BeninCity-AirportRoad	Edo	MPR	12.45	32.35	39.86	52.49	137.64	43.25	14.86	1.21	2.86
NIG14	Maiduguri-GRA	Borno	MPR	16.06	31.86	39.64	51.46	176.69	42.53	14.62	1.51	6.23
NIG02	Lagos-Lekki	Lagos	MPR	14.18	31.46	38.41	51.00	142.06	41.85	14.26	1.34	4.34
NIG04	Abuja-Asokoro	FCT	LPR	13.62	29.12	35.37	46.70	122.27	38.40	12.85	1.18	2.67
NIG11	Enugu-Coal Camp	Enugu	LPR	13.99	26.96	33.33	43.37	130.81	35.93	12.02	1.26	4.10
NIG15	Calabar-Marian	Cross River	LPR	11.69	25.16	31.13	40.79	103.96	33.74	11.46	1.22	3.08

Agglomerative Hierarchical Clustering

Ward's hierarchical clustering categorized 15 Nigerian monitoring stations into three distinct clusters: High Pollution Region (HPR), Medium Pollution Region (MPR), and Low Pollution Region (LPR), according to patterns of PM_{2.5} concentrations. This method aligns with hierarchical agglomerative clustering used in African air quality networks, successfully identifying geographically relevant pollution zones characterized by unique health exposure profiles (Izah et al., 2024). The dendrogram in figure 2 indicated optimal cluster separation at normalized linkage distances similar to

established thresholds in comparable studies. The concentration of HPR clusters in industrialized centers such as Kano and Port Harcourt facilitates targeted public health interventions for populations facing increased PM_{2.5}-related cardiorespiratory issues. Additionally, MPR and LPR classifications aid in the implementation of differentiated monitoring and policy protocols. The cluster-based framework converts spatially diverse pollution data into practical epidemiological insights that conform to risk-stratified air quality management standards.




Figure 2: A dendrogram illustrating the Hierarchical Clustering of 15 PM_{2.5} Monitoring Stations in Nigeria, Employing Ward's Method Alongside Euclidean Distance for the Years 2019 to 2023. The Red Dashed Line Signifies the Ideal cut-off for Three Distinct Clusters. HPR (High Pollution Regions), MPR (Medium Pollution Regions), and LPR (Low Pollution Regions)

High Pollution Regions (HPR)

Ward's clustering analysis in table 4 revealed three distinct PM_{2.5} exposure zones, each carrying important implications for health policy. The High Pollution Region (HPR), which encompasses 20% of monitoring stations, includes Kano-Industrial and Port Harcourt-Trans Amadi, both recognized as industrial hotspots. The mean PM_{2.5} concentrations in this area is 61.66 ± 21.69 µg m⁻³, exceeding Nigeria's national standard of 25 µg m⁻³ by a factor of 2.5 and surpassing WHO guidelines by a factor of 12.3 (World Health Organization, 2021). The concentration range recorded in Port Harcourt (32.3–184.0 µg m⁻³) reflects the episodic gas flaring and refinery emissions characteristic of the Niger Delta region. The Medium Pollution Region (MPR), encompassing nine significant urban centers and representing 60% of monitoring stations, exhibited an average particulate matter concentration of 46.69 ± 16.45 µg m⁻³. The current level is 8 -10 times above

the recommendations established by the World Health Organization, with vehicular emissions accounting for approximately 32% of the PM_{2.5} burden. The Low Pollution Region (LPR; 20% of stations) exhibited a mean concentration of 36.02 ± 12.27 µg m⁻³, exceeding the WHO annual guidelines by 7.2 times and surpassing WHO Interim Target-4 (35 µg m⁻³) (World Health Organization, 2021). This suggests that even the least polluted monitoring areas in Nigeria pose significant public health risks. This three-tier stratification identifies geographically distinct populations, necessitating proportionally scaled intervention strategies. Emission source controls are implemented in high pollution risk (HPR) industrial zones, while vehicle emission standards and traffic management are applied in moderate pollution risk (MPR) urban centers. Additionally, baseline monitoring continues in low pollution risk (LPR) areas.

Table 4: Summary Statistics by Region Type (2019-2023)

Region Type	NStations	Mean (µg/m ³)	PM _{2.5} SD (µg/m ³)	Min (µg/m ³)	Max (µg/m ³)	Median (µg/m ³)	Q1 (µg/m ³)	Q3 (µg/m ³)
HPR	3	61.66	21.69	18.64	241.32	56.86	45.78	74.63
MPR	9	46.69	16.45	12.45	189.00	43.21	34.76	56.39
LPR	3	36.02	12.27	11.69	130.81	33.41	26.96	43.36

Spatial Pattern Analysis

The spatial distribution map in Figure 3 illustrates clear geographical clustering patterns. Northern Nigeria, particularly Kano State, has been recognized as a notable pollution hotspot, with urban and industrial stations classified

as high pollution risk (HPR). The identified pattern reflects the characteristics of the area's semi-arid climate, which promotes dust resuspension, alongside vigorous commercial activities and a lack of vegetation to facilitate pollutant deposition.




Figure 3: Analysis of the Spatial Distribution of PM_{2.5} Monitoring Stations Throughout Nigeria Categorized by Pollution Levels. Red Circles Indicate Areas with High Levels of Pollution, Orange Circles Denote Regions with Medium Pollution Levels, and Green Circles Signify Areas with Low Pollution Levels. The Size of the Circle Indicates the Level of Pollution

The Niger Delta region, encompassing Rivers, Delta, and Edo states, exhibits varied patterns, featuring industrial stations in HPR and urban stations in MPR. This distribution highlights the significant impact of oil and gas operations on pollution levels in this region (Ukpong et al., 2019). Port Harcourt has been noted for its exceptionally high PM_{2.5} concentrations in sub-Saharan Africa, with measurements sometimes surpassing 500 µg m⁻³ during severe pollution events. Although Lagos is recognized as the largest metropolis in West Africa, its coastal position and the influence of prevailing winds contribute to a degree of natural dispersion of pollutants. Traffic congestion contributes to significant PM_{2.5} emissions from vehicular exhaust.

Temporal Annual Trends in PM_{2.5} Concentrations

The five-year trend analysis in Figure 4 and Table 5 reveals distinct temporal patterns with important implications for

public health concerning the three pollution clusters. The High Pollution Region (HPR) and Moderate Pollution Region (MPR) peaked in 2019, saw a notable decrease in 2020 owing to reduced industrial and vehicular activity during the COVID-19 lockdown, and reverted to near-baseline levels by 2023, underscoring the transient character of emission reductions in the absence of structural policy reforms. The Low Pollution Region (LPR) displayed relative stability alongside increasing variability, suggesting localized emissions linked to urban expansion, traffic growth, and industrial activities. The results demonstrate that even areas in Nigeria with lower pollution levels are affected by human activities, highlighting the need for continuous emission control strategies. The implementation of stricter vehicular standards, industrial monitoring, and strategic land-use planning is essential to mitigate long-term health risks associated with chronic PM_{2.5} exposure.




Figure 4: Analysis of Temporal Annual Trends in PM_{2.5} Concentrations Throughout Nigeria from 2019 to 2023. The Upper Panel Illustrates Monthly Fluctuations Characterized by Seasonal Trends, Whereas the Lower Panel Presents Yearly Averages Categorized by Region Type. Dashed Lines Represent the Interim Targets Set by the WHO

Table 5: Annual PM_{2.5} and AQI Extremes by Region Type

Region Type	Year	Mean PM _{2.5} (µg/m ³)	Max PM _{2.5} (µg/m ³)	Mean AQI	Max AQI	Max AQI Status
HPR	2019	64.76	197.70	148.84	273	Very Unhealthy
HPR	2020	58.47	241.32	141.92	312	Hazardous
HPR	2021	60.09	197.61	144.33	272	Very Unhealthy
HPR	2022	61.47	195.73	146.04	271	Very Unhealthy
HPR	2023	63.51	211.13	148.76	286	Very Unhealthy
LPR	2019	37.54	100.92	106.76	183	Unhealthy
LPR	2020	33.88	106.60	99.24	187	Unhealthy
LPR	2021	34.99	96.83	101.64	180	Unhealthy
LPR	2022	36.11	113.90	103.73	192	Unhealthy
LPR	2023	37.60	130.81	106.54	206	Very Unhealthy
MPR	2019	48.90	186.90	127.21	262	Very Unhealthy
MPR	2020	44.01	166.46	118.92	242	Very Unhealthy
MPR	2021	45.56	165.75	121.54	241	Very Unhealthy
MPR	2022	46.89	181.85	123.80	257	Very Unhealthy
MPR	2023	48.12	189.00	125.95	264	Very Unhealthy

Monthly and Seasonal Patterns

Figure 5 presents the monthly and seasonal fluctuations of PM_{2.5} concentrations across various regions. In the dry season (November–March), concentrations significantly increased, reaching their peak between December and February, coinciding with the Harmattan period. Dry-season levels were, on average, 35–58% higher than those observed during the rainy season (April–October). Mean monthly PM_{2.5} concentrations in the High Pollution Region (HPR) varied from 45 µg m⁻³ during July–August to 85 µg m⁻³ in January–February. In contrast, the Moderate Pollution Region (MPR) and Low Pollution Region (LPR) exhibited ranges of 35–62 µg m⁻³ and 28–48 µg m⁻³, respectively. The results align with findings from West African studies (e.g., Evans et al., 2022), which attribute 20–60% seasonal increases in PM_{2.5} to dust

intrusion and meteorological conditions during Harmattan, thereby reaffirming the significant seasonal impact on air quality dynamics in Nigeria.

Peak Concentrations and Timing

The Kano industrial station reported the highest PM_{2.5} concentration of 197.70 µg m⁻³ on March 12, 2019. Figure 5 and Table 6 show all locations experienced high amounts in September and October. September had an average daily HPR high of 142 µg m⁻³, with several days exceeding 150 µg. The MPR peaked at 186.90 µg m⁻³ in Warri-Effurun on October 31, 2019, due to biomass burning. Additionally, the LPR peaked at 100.92 µg m⁻³ in Abuja-Asokoro on October 10, 2019.



Figure 5: Analysis of Monthly PM_{2.5} Concentrations Throughout the 2019 Haze Episode in Three Distinct Pollution Regions. The Shaded Area (September–October) Represents the Peak Haze Period, Pollution from Local Emissions, Resulted in the Highest Recorded Concentrations

The observed values significantly surpassed the WHO's 24-hour guideline of 15 µg m⁻³, as well as the Nigerian national

standard of 50 µg m⁻³ (Air Quality Life Index, 2025; WHO, 2021).

Table 6: 2019 Haze Maximum Recorded Values by Region

Region Type	Date	Station Code	PM _{2.5} Max(µg/m ³)	AQI Max	Health Status	N Stations	Mean Annual PM _{2.5} (µg/m ³)
HPR	2019-03-12	NIG06	197.70	273	Very Unhealthy	3	64.76
MPR	2019-10-31	NIG13	186.90	262	Very Unhealthy	9	48.90
LPR	2019-10-10	NIG04	100.92	183	Unhealthy	3	37.54

Air Quality Index During Haze Episode

In Figure 6, the mean AQI for HPR was recorded at 148.84, with a peak maximum of 273, categorizing it as Very Unhealthy during the period. The distribution of AQI

categories revealed that Good (0-50) is 0%, Moderate (51-100) is 6.12%, Unhealthy for Sensitive Groups (101-150) is 36.26%, Unhealthy (151-200) is 55.62%, Very Unhealthy (201-300) is 2.01%, and Hazardous (301+) is 0%.




Figure 6: Analysis of the Distribution of Air Quality Index Categories Across Various Pollution Regions, Highlighting the Percentage of days within Each Health Category from 2019 to 2023. HPR Encountered Adverse Conditions on 52% of days, Whereas LPR also Faced Subpar Air Quality on more than Half of the Days. Throughout the Study Period, no Region Attained Good Air Quality at any Point

MPR showed a mean AQI of 127.21, peaking at 262, which is Very Unhealthy. Moderate, 24.84%. Unhealthy for Sensitive Groups, 41.67%. Unhealthy, 33.27%. Very Unhealthy, 0.21%. LPR's mean AQI was 106.76, with a maximum of 183, which is unhealthy. Moderate, 51.69%. Unhealthy for Sensitive Groups, 38.36%. Unhealthy, 9.95%. Every region shows Unhealthy levels during September and October, while HPR and MPR reached Very Unhealthy levels that required public health warnings.

AQI Distribution

Nigeria's five-year dataset showed sustained poor air quality in Figure 7. HPR's mean AQI was 146.4, indicating an unhealthy threshold for sensitive groups. The data shows that 94.8% of days had an AQI over 100 and 52.3% over 150, making them unhealthy. 3.1% of days had an AQI over 200, which is Very Unhealthy.




Figure 7: Comparative Boxplots Illustrating the Distributions of PM_{2.5} Concentrations (Left) and Air Quality Index Values (Right) Across the Three Pollution Regions from 2019 to 2023. Reference Lines Denote Established Guidelines and Air Quality Index Thresholds

The MPR showed a mean AQI of 123.7, an unhealthy level for sensitive groups. The data showed that 76.2% of days had an AQI above 100, 28.4% above 150, and 0.8% above 200. The LPR had an unhealthy for sensitive groups mean AQI of 103.6. The data shows that 54.6% of days had an AQI above 100, 8.2% above 150, and 0.3% above 200. In Nigeria's least

polluted areas, air quality standards were surpassed on more than half of the observed days. These conditions pose ongoing health risks for children, the elderly, and individuals with respiratory or cardiovascular issues (US EPA, 2024).

Health Impact Assessment in Global Context

Nigerian residents face an increasing health burden across pollution zones, as classified by the US EPA AQI. Residents of HPR encounter "unhealthy" air quality conditions (AQI 151–200), which are associated with increased risks of respiratory dysfunction and cardiovascular stress in the general population. In contrast, MPR residents experience "unhealthy for sensitive groups" conditions (AQI 101–150), impacting 30% of exposed children and adults who exhibit moderate or greater lung function impairment during moderate exertion. LPR regions surpass an AQI of 100 in 55% of measurements, resulting in all exposure zones exceeding internationally accepted safety thresholds. Examining these levels in relation to regional and global datasets highlights the seriousness of Nigeria's pollution issue. Nigerian HPR concentrations ($61.66 \mu\text{g m}^{-3}$) surpass those of significant West African counterparts Accra, Ghana ($49.5 \mu\text{g m}^{-3}$) and Lomé, Togo ($23.5 \mu\text{g m}^{-3}$)—and are comparable to Delhi, India ($92.7 \mu\text{g m}^{-3}$). In contrast, Nigerian LPR ($36.02 \mu\text{g m}^{-3}$) is similar to levels in Beijing, China ($32.6 \mu\text{g m}^{-3}$) and Bangkok, Thailand ($21.7 \mu\text{g m}^{-3}$), suggesting that even Nigeria's least-polluted stations exceed urban baselines in developed nations (US urban average: 7–12 $\mu\text{g m}^{-3}$). This gradient illustrates that Nigeria's air quality crisis is pervasive throughout its monitored regions, indicating the need for immediate structural emission interventions instead of sporadic policy measures.

CONCLUSION

In this study, a comprehensive nationwide and regionally comparative classification of PM_{2.5} pollution in Nigeria was explored, utilizing agglomerative hierarchical clustering (AHC) with Ward's linkage. This study examined five years of continuous observations (2019–2023) from 15 monitoring stations, categorizing the data into three distinct zones: High Pollution Region (HPR), Medium Pollution Region (MPR), and Low Pollution Region (LPR). These zones collectively illustrate the diverse industrial, urban, and climatic characteristics of Nigeria. The findings reveal a distinct pollution gradient, defined by HPR zones (e.g., Kano and Port Harcourt) predominantly impacted by industrial and combustion-related emissions, MPR zones influenced by considerable urbanization and vehicular activity, and LPR zones exhibiting comparatively lower, yet still notable, particulate concentrations pertinent to health. This hierarchical classification offers a comprehensive spatial overview of air quality in Nigeria, transcending analyses centered on individual monitoring stations and enhancing the understanding of spatial variations in pollutant exposure.

The approach attempts to demonstrate robustness. However, limitations remain due to inconsistent station coverage and reliance on a single clustering algorithm. To overcome these limitations, integrating satellite-derived PM_{2.5} data is crucial for enhancing spatial completeness, alongside the adoption of hybrid machine learning or source apportionment models to improve clustering accuracy. Targeted mitigation strategies are essential from a policy standpoint, encompassing the regulation of industrial emissions and enforcement of fuel standards in HPR zones, improved urban transport management in MPR zones, and proactive land-use and agricultural controls in LPR zones. Future research should build upon this framework by incorporating meteorological and chemical composition data to strengthen the link between emission sources and observed pollution patterns, thus supporting evidence-based air quality management and public health protection in Nigeria.

REFERENCES

- Abayomi, O., Olayemi, T. E., & Ogunbade, T. (2021). Environmental pollution and its ecological consequences on the Niger Delta: A review of the literature. *African Journal of Environment and Natural Science Research*, 4, 27-42.
- Abuh, P. O., & John, J. A. (2023). Geographic Analysis of Traffic Congestion in FCT-Abuja, Nigeria. *Journal of Sustainability and Environmental Management*, 2(1), 26-32.
- Abulude, F. O., Oyetunde, J. G., & Feyisetan, A. O. (2024). Air Pollution in Nigeria: A Review of Causes, Effects, and Mitigation Strategies. *Continental Journal of Applied Sciences*, 19, 1-23.
- Adeleye, M. A., Connor, S. E., & Haberle, S. G. (2021). A quantitative synthesis of Holocene vegetation change in Nigeria (Western Africa). *The Holocene*, 31(11-12), 1681-1689.
- Almsatar, T. (2020). Environmental issues of biomass-burning in sub-Saharan African countries. In *Biomass burning in sub-Saharan Africa: Chemical issues and action outreach* (pp. 1-14). Dordrecht: Springer Netherlands.
- Arunachalam, G. (2024). A Human Health Risk Assessment (HHRA) associated with gaseous pollutants in the ambient environment at Coimbatore City's High Traffic Road.
- Azizan, N. A., Othman, A. S., Meramat, A. A., Amin, S. N. S. M., & Azid, A. (2023). A framework to spatially cluster air quality monitoring stations in Peninsular Malaysia using the hybrid clustering method. *Malaysian Journal of Fundamental and Applied Sciences*, 19(5), 804-816.
- Balaji, R., Bapat, R. B., & Goel, S. (2022). Generalized Euclidean distance matrices. *Linear and Multilinear Algebra*, 70(21), 6908-6929.
- Benabed, A., & Boulbair, A. (2022). PM₁₀, PM_{2.5}, PM₁, and PM_{0.1} resuspension due to human walking. *Air Quality, Atmosphere & Health*, 15(9), 1547-1556.
- Britannica, T. Editors of Encyclopaedia (2025). List of cities and towns in Nigeria. <https://www.britannica.com/topic/list-of-cities-and-towns-in-Nigeria-2042462>. Accessed 23rd October, 2025
- Busolo, W. S., & Njabira, V. I. (2022). Air Quality. In *The Palgrave Handbook of Urban Development Planning in Africa* (pp. 327-372). Cham: Springer International Publishing.
- El-Araby, I. E., Moawed, S. A., Hassan, F. A., & Gouda, H. F. (2023). A combined approach of multiple correspondence analysis and hierarchical cluster analysis for profiling relationships among categorical risk predictors: a bluetongue case study. *Slovenian Veterinary Research/Slovenski Veterinarski Zbornik*, 60.
- Elechi, F. A., Etea, E. A., & Mbanusi, E. C. (2023). Governance, security and economic cooperation in the southeast geo-political zone, Nigeria. *South east Journal of Political Science*, 9(1).
- Evans, M. J., et al. (2022). A network of field-calibrated low-cost sensor measurements of PM_{2.5} in West Africa. *ACS*

- Earth and Space Chemistry*, 6(4), 1048-1062. <https://doi.org/10.1021/acsearthspacechem.1c00391>
- Ezeh, G. C., Obioh, I. B., Asubiojo, O., Abiye, O. E., & Onyeuwaoma, N. D. (2019). A study of PM_{2.5}–10 pollution at three functional receptor sites in a sub-Saharan African megacity. *Aerosol Science and Engineering*, 3(3), 65-74.
- Fu, D., Song, Z., Zhang, X., Wu, Y., Duan, M., Pu, W., ... & Xia, X. (2020). Similarities and Differences in the Temporal Variability of PM_{2.5} and AOD between Urban and Rural Stations in Beijing. *Remote Sensing*, 12(7), 1193.
- Garcia, A., Santa-Helena, E., De Falco, A., de Paula Ribeiro, J., Gioda, A., & Gioda, C. R. (2023). Toxicological effects of fine particulate matter (PM_{2.5}): health risks and associated systemic injuries—systematic review. *Water, Air, & Soil Pollution*, 234(6), 346.
- Glenn, B. E., Espira, L. M., Larson, M. C., & Larson, P. S. (2022). Ambient air pollution and non-communicable respiratory illness in sub-Saharan Africa: a systematic review of the literature. *Environmental Health*, 21(1), 40.
- Horn, S. A., & Dasgupta, P. K. (2024). The Air Quality Index (AQI) in historical and analytical perspective a tutorial review. *Talanta*, 267, 125260.
- Ifie, I., Igwebuikwe, C. G., Imasuen, P., Akalamudo, W., Ogheneborhie, O., Akpodiete, J. O., & Eze, U. A. (2022). Assessment of aflatoxin and heavy metals levels in maize and poultry feeds from Delta State, Nigeria. *International Journal of Environmental Science and Technology*, 19(12), 12551-12560.
- Izah, S. C., Sylva, L., Ogwu, M. C., Shahsavani, A., Bazzazpour, S., & Rahmatinia, M. (2024). Modeling and statistical approaches for air pollution analysis. In *Air Pollutants in the Context of One Health: Fundamentals, Sources, and Impacts* (pp. 425-456). Cham: Springer Nature Switzerland.
- Jalali, S., Karbakhsh, M., Momeni, M., Taheri, M., Amini, S., Mansourian, M., & Sarrafzadegan, N. (2021). Long-term exposure to PM_{2.5} and cardiovascular disease incidence and mortality in an Eastern Mediterranean country: findings based on a 15-year cohort study.
- Jin, X., Fiore, A. M., Civerolo, K., Bi, J., Liu, Y., Van Donkelaar, A., ... & Kinney, P. L. (2019). Comparison of multiple PM_{2.5} exposure products for estimating health benefits of emission controls over New York State, USA. *Environmental Research Letters*, 14(8), 084023.
- Khomenko, S., Cirach, M., Pereira-Barboza, E., Mueller, N., Barrera-Gómez, J., Rojas-Rueda, D., ... & Nieuwenhuijsen, M. (2021). Premature mortality due to air pollution in European cities: a health impact assessment. *The Lancet Planetary Health*, 5(3), e121-e134.
- Kodiya, M. A., Modu, M. A., Ishaq, K., Yusuf, Z., Wakili, A. Z., Dayyabu, N., ... & Babangida, M. U. (2025). Environmental pollution in Nigeria: Unlocking integrated strategies for environmental sustainability. *African Journal of Environmental Sciences and Renewable Energy*, 18(1), 30-50.
- Lawal, H., & Muhammed, M. I. (2022). Investigating the Annual Atmospheric Pollution and Its Analysis. *FUDMA JOURNAL OF SCIENCES*, 6(5), 102-108.
- Léon, J. F., Akpo, A. B., Bedou, M., Djossou, J., Bodjrenou, M., Yoboué, V., & Liousse, C. (2021). PM_{2.5} surface concentrations in southern West African urban areas based on sun photometer and satellite observations. *Atmospheric Chemistry and Physics*, 21(3), 1815-1834.
- Magi, B. I., Cupini, C., Francis, J., Green, M., & Hauser, C. (2020). Evaluation of PM_{2.5} measured in an urban setting using a low-cost optical particle counter and a Federal Equivalent Method Beta Attenuation Monitor. *Aerosol Science and Technology*, 54(2), 147-159.
- Maldonado-Salguero, P., Bueso-Sánchez, M. C., Molina-García, Á., & Sánchez-Lozano, J. M. (2022). Spatio-temporal dynamic clustering modeling for solar irradiance resource assessment. *Renewable Energy*, 200, 344-359.
- Manisalidis, I., Stavropoulou, E., Stavropoulos, A., & Bezirtzoglou, E. (2020). Environmental and health impacts of air pollution: a review. *Frontiers in public health*, 8, 14.
- Meo, S. A., Salih, M. A., Alkhalifah, J. M., Alsomali, A. H., & Almushawah, A. A. (2024). Environmental pollutants particulate matter (PM_{2.5}, PM₁₀), Carbon Monoxide (CO), Nitrogen dioxide (NO₂), Sulfur dioxide (SO₂), and Ozone (O₃) impact on lung functions. *Journal of King Saud University-Science*, 36(7), 103280.
- Mitchell, L., & Bala, I. (2024). Thematic exploration of educational research after the COVID pandemic through topic modelling. *Journal of Applied Learning & Teaching*, 7(1), 26-34.
- Nnadi, V. E., Udokporo, E. L., & Okolo, O. J. (2022). Petroleum production activities and depletion of biodiversity: A case of oil spillage in the Niger Delta. In *Handbook of Environmentally Conscious Manufacturing* (pp. 95-111). Cham: Springer International Publishing.
- Ogwu, M. C., Imarhiagbe, O., Ikhajiagbe, B., & Osawaru, M. E. (2024). Toward understanding the impacts of air pollution. In *Sustainable Strategies for Air Pollution Mitigation: Development, Economics, and Technologies* (pp. 3-43). Cham: Springer Nature Switzerland.
- Oluwakoya, A. O. (2024). A comprehensive assessment of transportation emissions in Nigeria: Trends, drivers, and impacts. *Proceedings of the Nigerian Academy of Science*, 16(1s), 61-71.
- Omokaro, G. O., Efeni, O. S., Adeyanju, O. I., Obomejoro, J., & Akpotu, E. (2025). Oil spillage in the Niger Delta: impacts, institutional failures, and policy reforms. *Academia Environmental Sciences and Sustainability*, 2(3).
- Paparrizos, J., Yang, F., & Li, H. (2024). Bridging the gap: A decade review of time-series clustering methods. *arXiv preprint arXiv:2412.20582*.
- Pyae, T. S. (2021). Temporal distribution of air pollutants (PM_{2.5}, PM₁₀ and O₃) in Yangon City, Myanmar during 2019-2021.

- Raheja, G., Sabi, K., Sonla, H., Gbedjangni, E. K., McFarlane, C. M., Hodoli, C. G., & Westervelt, D. M. (2022). A network of field-calibrated low-cost sensor measurements of PM_{2.5} in Lomé, Togo, over one to two years. *ACS Earth and Space Chemistry*, 6(4), 1011-1021.
- Rahman, E. A., Hamzah, F. M., Latif, M. T., & Dominick, D. (2022). Assessment of PM_{2.5} patterns in Malaysia using the clustering method. *Aerosol and Air Quality Research*, 22(1), 210161.
- Roy, A., Mostafa, M. G., & Saha, M. K. (2025). Air pollutant dynamics and behaviours in tobacco processing and storage environments: implications for air quality and health hazards. *Environmental Science: Atmospheres*.
- Singh, S., Kulshrestha, M. J., Rani, N., Kumar, K., Sharma, C., & Aswal, D. K. (2023). An overview of vehicular emission standards. *Mapan*, 38(1), 241-263.
- Sooktawee, S., Kanchanasuta, S., & Bunplod, N. (2023). Assessment of 24-h moving average PM_{2.5} concentrations in Bangkok, Thailand against WHO guidelines. *Sustainable Environment Research*, 33(1), 3.
- Sufiyan, I., Mohammed, K. D., Bello, I. E., & Zaharadeen, I. (2020). Impact of harmattan season on human health in Keffi, Nasarawa State, Nigeria. *Matrix Science Medica*, 4(2), 44-50.
- Sunday, I. (2022). Assessment of Air Quality Across Different Land Uses in Gwagwalada Town, FCT-Abuja, Nigeria. *Fudma Journal of Sciences*.
- Ukpong, I. G., Balcombe, K. G., Fraser, I. M., & Areal, F. J. (2019). Preferences for mitigation of the negative impacts of the oil and gas industry in the Niger Delta region of Nigeria. *Environmental and Resource Economics*, 74(2), 811-843.
- US EPA. (2024). *Final updates to the Air Quality Index (AQI) for particulate matter*. EPA Fact Sheet. Retrieved October 20, 2025, from <https://www.epa.gov/system/files/documents/2024-02/pm-naaqs-air-quality-index-fact-sheet.pdf>
- Wambebe, N. M., & Duan, X. (2020). Air quality levels and health risk assessment of particulate matters in Abuja municipal area, Nigeria. *Atmosphere*, 11(8), 817.
- World Health Organization. (2021). *WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide*. World Health Organization.
- World Health Organization. (2022). *Setting global research priorities for urban health*. World Health Organization.
- Xu, J., Xu, M., Zhao, Y., Wang, S., Tao, M., & Wang, Y. (2021). Spatial-temporal distribution and evolutionary characteristics of water environment sudden pollution incidents in China from 2006 to 2018. *Science of the Total Environment*, 801, 149677.
- Zhou, B., Lin, K., Quan, J., Liao, J., Chen, Y., & Xu, X. (2025). Comprehensive insights into the disease burden linked to air pollution exposure across US states: Findings from the 2021 global burden of disease study. *Frontiers in Public Health*, 13, 1636544.



©2025 This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International license viewed via <https://creativecommons.org/licenses/by/4.0/> which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is cited appropriately.